



# New Frontiers in Early-Universe Cosmology

**Anna Ijjas**

Max Planck Institute for Gravitational Physics  
(Albert Einstein Institute)



characteristic features of the very early universe



As we make progress in our understanding,  
our view of which problems are ‘hard’ changes.

## ‘hard’ problems:

- singularity resolution & its cosmological implications  
Brandenberger; Graham, Kaplan, Rajendran; Ijjas & Steinhardt; Vafa, ...
- models based on modified gravity  
Ijjas, Steinhardt, Pretorius; Lehner, Kovacs & Reall, ...
- non-linear & non-perturbative cosmological dynamics  
within Einstein gravity  
Clough, Flauger; East, Senatore; Ijjas, Steinhardt, Pretorius

# Cosmology & General Relativity

$$R_{\alpha\beta} = T_{\alpha\beta} - \frac{1}{2}g_{\alpha\beta}T_{\lambda}^{\lambda}$$

?



$$\nabla^{\beta}T_{\alpha\beta} = 0$$

$$H^{-1} = (a/a_0)^{\varepsilon}$$

$$\varepsilon = -\dot{H}/H^2$$

# conventional picture of smoothing & flattening:

Step #1: a Hubble patch becomes smooth following a ‘natural succession’ of events

$$H^2 = \left( \frac{\dot{a}}{a} \right)^2 = \frac{1}{3} \left( \frac{\rho_m^0}{a^3} + \frac{\rho_r^0}{a^4} \right) - \frac{k}{a^2} + \frac{\sigma^2}{a^6} + \frac{1}{3} \left( \frac{\rho_\phi^0}{a^{2\epsilon}} \right)$$

“survival of the largest”

Step #2: space-time becomes flat and smooth on super-Hubble scales while the Hubble radius  $|H|^{-1}$  evolves at a different rate than the scale factor  $a$ :

$$|H|^{-1} \sim a^\epsilon$$


‘rate’ is determined by the stress-energy content characterized by the

equation-of-state  $\epsilon \equiv \frac{3}{2} \left( 1 + \frac{p}{\varrho} \right)$

What can cause slow contraction ( $\varepsilon \gg 3$ )?

$$\varepsilon = \frac{3\left(\frac{1}{2}\dot{\phi}^2\right)}{\frac{1}{2}\dot{\phi}^2 + V(\phi)}$$

$$V < 0 \Rightarrow \varepsilon > 3$$




# slow contraction is different!

w/W.G. Cook, E. Y. Davies, D. Garfinkle, I. Glushchenko, F. Pretorius, P.J. Steinhardt, and A.P. Sullivan

arXiv: 2109.09768 ; 2104.12293; 2103.00584; 2006.04999; 2006.01172


$$\begin{array}{ccc} R_{\alpha\beta} = T_{\alpha\beta} - \frac{1}{2}g_{\alpha\beta}T_{\lambda}^{\lambda} & \xrightarrow{\text{?}} & H^{-1} = (a/a_0)^{\varepsilon} \\ \nabla^{\beta}T_{\alpha\beta} = 0 & & \varepsilon = -\dot{H}/H^2 \end{array}$$



# Smoothing through Ultralocality

arXiv: 2104.12293; 2103.00584; 2006.04999; 2006.01172

$$\partial_t \bar{n}_{ab} = -\left(\mathcal{N} - 1\right) \bar{n}_{ab} + \mathcal{N} \left( 2\bar{n}_{(a}{}^c \bar{\Sigma}_{b)c} - \epsilon^{cd} {}_{(a} \bar{E}_c{}^i \partial_i \bar{\Sigma}_{b)d} \right) - \epsilon^{cd} {}_{(a} \bar{\Sigma}_{b)d} \bar{E}_c{}^i \partial_i \mathcal{N}$$



# Generation of primordial perturbations

e.g., Ijjas et al.: PRD 89 (2014)123520


$$\mathcal{S} = \int d^4x \sqrt{-g} \left( \frac{1}{2}R - \frac{1}{2}(\partial_\mu\phi)^2 + V_0 \exp(-\sqrt{2\epsilon}\phi) - \frac{1}{2}\Omega^2(\phi)(\partial_\mu\chi)^2 \right)$$

FRW BACKGROUND:

$$H^2 = \frac{1}{3} \left( \frac{1}{2}\dot{\phi}^2 + \frac{1}{2}\Omega^2(\phi)\dot{\chi}^2 + V(\phi) \right),$$



$$\ddot{\phi} + 3H\dot{\phi} - \Omega\Omega_{,\phi}\dot{\chi}^2 + V_{,\phi} = 0,$$

$$\ddot{\chi} + \left( 3H + 2\frac{\dot{\Omega}}{\Omega} \right) \dot{\chi} = 0,$$



# Non-perturbative analysis

arXiv: 2109.09768



Time=0

time evolution of the entropic field's  
kinetic energy density

Time=0

time evolution of the background field

# Non-perturbative analysis

arXiv: 2109.09768

**shear - blue**  
**curvature - red**  
**matter - green**

